

# Wavelet Approximations of Boundary Value Problems

Consider the problem

$$\begin{cases} Au = f & \text{in } \Omega \\ Bu = 0 & \text{on } \partial\Omega . \end{cases}$$

where  $A, B$  are linear differential operators.

Assume that the problem can be formulated as:

*Given  $f \in V'$ , find  $u \in V$  such that*

$$\langle Au, v \rangle = \langle f, v \rangle, \quad \forall v \in V,$$

where

- $V = V^{(r)}$  is a Hilbert space of regularity  $r > 0$ , continuously embedded in  $L^2(\Omega)$ , which may incorporate boundary conditions
- $V'$  is the dual space of  $V$
- $\langle \cdot, \cdot \rangle$  is the duality pairing between  $V'$  and  $V$ .

## **Assumption:**

$A : V \rightarrow V'$  is an algebraic and topological isomorphism, so that

$$\|Av\|_{V'} \asymp \|v\|_V, \quad \forall v \in V.$$

This holds, e.g., if  $A$  is continuous from  $V$  to  $V'$  and coercive on  $V$ , i.e., there exists  $\alpha > 0$  such that

$$\langle Av, v \rangle \geq \alpha \|v\|_V^2, \quad \forall v \in V.$$

## Examples

### 1. Dirichlet Problem for the Poisson Equation

$$\begin{cases} -\Delta u = f & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega . \end{cases}$$

Set  $V = H_0^1(\Omega) = \{v \in H^1(\Omega) : v = 0 \text{ on } \partial\Omega\}$  equipped with the norm

$$\|v\|_{H_0^1(\Omega)} = \left( \int_{\Omega} |\nabla v|^2 dx \right)^{1/2} .$$

Then,

$$\langle Au, v \rangle := \int_{\Omega} \nabla u \cdot \nabla v dx, \quad \langle f, v \rangle := \int_{\Omega} f v dx$$

with  $f \in L^2(\Omega)$ .

### 2. Neumann Problem for the Helmholtz Equation

$$\begin{cases} -\Delta u + u = f & \text{in } \Omega \\ \frac{\partial u}{\partial n} = 0 & \text{on } \partial\Omega . \end{cases}$$

Set  $V = H^1(\Omega)$  equipped with the norm

$$\|v\|_{H^1(\Omega)} = \left( \int_{\Omega} |\nabla v|^2 dx + \int_{\Omega} |v|^2 dx \right)^{1/2} .$$

Then,

$$\langle Au, v \rangle := \int_{\Omega} \nabla u \cdot \nabla v dx + \int_{\Omega} u v dx, \quad \langle f, v \rangle := \int_{\Omega} f v dx .$$

Let us introduce a multilevel (wavelet) basis in  $V$ :

$$V = \text{span} \{ \psi_\lambda : \lambda \in \mathcal{M} \}$$

and let us set

$$u = \sum_{\lambda \in \mathcal{M}} u_\lambda \psi_\lambda, \quad \mathbf{u} = (u_\lambda)$$

$$a_{\mu\lambda} = \langle A\psi_\lambda, \psi_\mu \rangle, \quad \mathbf{A} = (a_{\mu\lambda})$$

$$f_\mu = \langle f, \psi_\mu \rangle, \quad \mathbf{f} = (f_\mu).$$


Then, the operator equation can be written as the (infinite order) algebraic system

$$\mathbf{A}\mathbf{u} = \mathbf{f}$$

Ordering the hierarchical basis functions by levels

$$(\psi_\lambda) = (\psi_{jk}) = ((\psi_{jk}), k \in \mathcal{K}_j), j \geq j_0,$$

the matrix  $\mathbf{A}$  has a typical non-banded, sparse block structure:



(compare with the banded matrix of a non-hierarchical basis)

## Discrete representations of Operators

Let  $V = V^{(r)}$  be a Hilbert space of regularity  $r$  and let  $\Psi = \{\psi_\lambda\}_{\lambda \in \mathcal{M}}$  be a Riesz basis in  $V$ :

$$\|v\|_V \asymp \|\mathbf{v}\|_{\ell^2(\mathcal{M})}, \quad \forall v = \sum_{\lambda \in \mathcal{M}} v_\lambda \psi_\lambda =: \mathbf{v}^T \Psi \in V.$$

Similarly, let  $W = W^{(s)}$  be a Hilbert space of regularity  $s$  and let  $\tilde{\Psi} = \{\tilde{\psi}_\lambda\}_{\lambda \in \mathcal{M}}$  be a Riesz basis in  $W$ .

Let  $A : V \rightarrow W$  be an algebraic and topological isomorphism,

$$\|Av\|_W \asymp \|v\|_V, \quad \forall v \in V.$$

For any  $\lambda \in \mathcal{M}$ , let

$$A\psi_\lambda =: \sum_{\mu \in \mathcal{M}} a_{\mu\lambda} \tilde{\psi}_\mu$$

be the expansion of  $A\psi_\lambda \in W$  according to the Riesz basis  $\tilde{\Psi}$ .

Set

$$\mathbf{A} = (a_{\mu\lambda})_{\mu, \lambda \in \mathcal{M}}$$

Then,

$$v = \mathbf{v}^T \Psi \in V \quad \text{iff} \quad Av = (\mathbf{A}\mathbf{v})^T \tilde{\Psi} \in W$$

and

$$\mathbf{A} : \ell^2(\mathcal{M}) \rightarrow \ell^2(\mathcal{M})$$

is an algebraic and topological isomorphism.

## Finite Dimensional (Galerkin) Approximation

Choose a finite subset  $\Lambda \subset \mathcal{M}$ . Set  $N = \text{card } \Lambda$  and

$$V_\Lambda = \text{span} \{ \psi_\lambda : \lambda \in \Lambda \}$$

Consider the Galerkin approximation

*Find  $u_\Lambda \in V_\Lambda$  such that*

$$\langle Au_\Lambda, v_\Lambda \rangle = \langle f, v_\Lambda \rangle, \quad \forall v_\Lambda \in V_\Lambda,$$

equivalent to the  $N \times N$  algebraic system

$$\mathbf{A}_\Lambda \mathbf{u}_\Lambda = \mathbf{f}_\Lambda$$

with

$$\mathbf{A}_\Lambda = (a_{\mu\lambda})_{\mu, \lambda \in \Lambda}$$

$$\mathbf{u}_\Lambda = (u_{\Lambda, \lambda})_{\lambda \in \Lambda} \quad \text{if} \quad u_\Lambda = \sum_{\lambda \in \Lambda} u_{\Lambda, \lambda} \psi_\lambda$$

$$\mathbf{f}_\Lambda = (f_\mu)_{\mu \in \Lambda}.$$

The multilevel structure of the basis can be exploited in

- preconditioning
- compression
- adaptivity

## Preconditioning

Basic facts from linear algebra:

Let  $\mathbf{A}$  be a  $N \times N$  non-singular matrix. Define

$$\text{cond}_2(\mathbf{A}) := \|\mathbf{A}\|_2 \|\mathbf{A}^{-1}\|_2.$$

Assume that  $\mathbf{A}$  is symmetric positive-definite, with eigenvalues  $\lambda > 0$ .

Then

$$\text{cond}_2(\mathbf{A}) = \frac{\lambda_{\max}}{\lambda_{\min}}.$$

Iterative solution of

$$\mathbf{Ax} = \mathbf{b}$$

by Richardson iterations

$$\mathbf{x}^{k+1} = \mathbf{x}^k + \alpha(\mathbf{b} - \mathbf{Ax}^k), \quad k = 0, 1, \dots$$

for some  $\alpha > 0$ .

The error  $\mathbf{e}^k = \mathbf{x}^k - \mathbf{x}$  satisfies

$$\mathbf{e}^{k+1} = (\mathbf{I} - \alpha\mathbf{A})\mathbf{e}^k,$$

whence

$$\|\mathbf{e}^k\|_2 \leq \rho^k \|\mathbf{e}^0\|_2$$

with

$$\rho := \rho(\mathbf{I} - \alpha\mathbf{A}) < 1 \quad \text{provided} \quad \alpha < \frac{2}{\lambda_{\max}}.$$

The factor  $\rho$  is minimized for

$$\alpha_{\text{opt}} := \frac{2}{\lambda_{\max} + \lambda_{\min}},$$

for which

$$\rho_{\text{opt}} = \frac{\lambda_{\max} - \lambda_{\min}}{\lambda_{\max} + \lambda_{\min}} = \frac{\text{cond}_2(\mathbf{A}) - 1}{\text{cond}_2(\mathbf{A}) + 1}$$

If  $\text{cond}_2(\mathbf{A}) \gg 1$ , the convergence is slow.

*Remedy:* Preconditioning. Replace

$$\mathbf{A}\mathbf{x} = \mathbf{b}$$

by

$$\mathbf{P}^T \mathbf{A}\mathbf{x} = \mathbf{P}^T \mathbf{b}$$

i.e.,

$$\mathbf{P}^T \mathbf{A}\mathbf{P}\mathbf{y} = \mathbf{P}^T \mathbf{b}, \quad \mathbf{y} = \mathbf{P}^{-1}\mathbf{x},$$

where the matrix  $\mathbf{P}$  is chosen in such a way that the symmetric matrix

$$\mathbf{B} := \mathbf{P}^T \mathbf{A}\mathbf{P}$$

satisfies

$$\text{cond}_2(\mathbf{B}) \ll \text{cond}_2(\mathbf{A}).$$

Preconditioning is easily accomplished by wavelets.

[Jaffard '92, Dahmen - Kunoth '92, Masson '98]

Assume that

- $A$  is a symmetric, continuous and coercive operator on  $V$ .
- The wavelet basis  $\{\psi_\lambda\}_{\lambda \in \mathcal{M}}$  is normalized in  $L^2(\Omega)$

**Theorem.**  $\mathbf{A}_\Lambda$  is spectrally equivalent to the diagonal matrix

$$\mathbf{D}_\Lambda = \text{diag} (2^{2r|\lambda|})_{\lambda \in \Lambda} .$$

Indeed, for any  $\mathbf{v}_\Lambda = (v_\lambda)_{\lambda \in \Lambda}$ , set  $v = \sum_{\lambda \in \Lambda} v_\lambda \psi_\lambda \in V$ . Then

$$\begin{aligned} \mathbf{v}_\Lambda^T \mathbf{A}_\Lambda \mathbf{v}_\Lambda &= \langle Av, v \rangle \asymp \|v\|_V^2 \\ &\asymp \sum_{\lambda \in \Lambda} 2^{2r|\lambda|} |v_\lambda|^2 = \mathbf{v}_\Lambda^T \mathbf{D}_\Lambda \mathbf{v}_\Lambda . \end{aligned}$$

**Corollary.** Set

$$\mathbf{B}_\Lambda = \mathbf{D}_\Lambda^{-1/2} \mathbf{A}_\Lambda \mathbf{D}_\Lambda^{-1/2}$$

Then,

$$\text{cond}_2(\mathbf{B}_\Lambda) \asymp 1$$

for all  $\Lambda \subset \mathcal{M}$ .

Equivalently, normalize the wavelets in  $V$  rather than in  $L^2(\Omega)$ :

$$\psi_\lambda^* := 2^{-r|\lambda|} \psi_\lambda .$$

Then,  $\mathbf{B}_\Lambda = (\langle A\psi_\lambda^*, \psi_\mu^* \rangle)_{\mu, \lambda \in \Lambda}$ .

For any  $\mathbf{v}_\Lambda = (v_\lambda)_{\lambda \in \Lambda}$ , set  $v^* = \sum_{\lambda \in \Lambda} v_\lambda \psi_\lambda^*$ . One has

$$\|v^*\|_V \asymp \|\mathbf{v}_\Lambda\|_{\ell^2(\Lambda)} \quad \text{and,} \quad \mathbf{v}_\Lambda^T \mathbf{B}_\Lambda \mathbf{v}_\Lambda \asymp \|\mathbf{v}_\Lambda\|_{\ell^2(\Lambda)}^2 .$$

*Improvements:*

- take  $D_\Lambda = \text{diag } A_\Lambda$  [Masson '97]
- Exploit the possibility of compressing the inverse  $A_\Lambda^{-1}$  using as  $D_\Lambda$  a 'Sparse Approximate Inverse' of  $A_\Lambda$  [Masson '97, Chan - Tang - Wan '97]

Remark:

Multilevel preconditioning is achieved also in other frameworks, such as:

- multigrid methods [Brandt '77, Hackbusch '85]
- hierarchical finite elements [Yserentant '86, Bank '88]
- BPX-type preconditioners [Bramble - Pasciak - Xu '90]

Set

$$P_j v = \sum_{k \in \mathcal{K}_j} \frac{(v, \Phi_{jk})_{L^2(\Omega)}}{(1, \Phi_{jk})_{L^2(\Omega)}} \Phi_{jk}$$

with  $\{\Phi_{jk}\}_{k \in \mathcal{K}_j}$  is the nodal (Lagrange) basis related to a finite element mesh of level  $j$

Setting

$$Q_j v = P_{j+1} v - P_j v$$

one has

$$\|v\|_{H^1(\Omega)} \sim \sum_{j \geq j_0} 2^{2j} \|Q_j v\|_{L^2(\Omega)}^2, \quad \forall v \in H^1(\Omega).$$

## Compression

[Beylkin - Coifman - Rokhlin '91, Dahmen - Prössdorf -Schneider '93, von Petersdorf - Schwab '95]

Many non-zero entries  $a_{\mu\lambda}$  of the matrix  $\mathbf{A}$  are indeed small, if level difference  $||\lambda| - |\mu||$  is large, due to the moment condition.

Consider, for instance

$$a_{\mu\lambda} = \int_0^1 \psi'_{jk}(x) \psi'_{\ell h}(x) dx.$$

Then

$$|a_{\mu\lambda}| \lesssim 2^{j+\ell} 2^{-\rho|j-\ell|} \chi_{\text{supp}\psi_\lambda \cap \text{supp}\psi_\mu}$$

where  $\rho = \min(\tilde{L} + 1, \sigma - 1)$

Indeed, assuming e.g.  $j \leq \ell$ ,

$$\begin{aligned} a_{\mu\lambda} &= 2^{j+\ell} 2^{(j-\ell)/2} \int_0^1 \psi'(2^{j-\ell}y - m) \psi'(y) dy \\ &= 2^{j+\ell} 2^{(j-\ell)/2} \int_0^1 [\psi'(2^{j-\ell}y - m) - p(y)] \psi'(y) dy \end{aligned}$$

for some  $m$  and for any polynomial  $p$  of degree  $\leq \tilde{L}$ .

By Whitney's Theorem (recall Jackson inequality) one has

$$\begin{aligned} \|\psi'(2^{j-\ell} \cdot - m) - p\|_{L^2(0,1)} &\lesssim |\psi'(2^{j-\ell} \cdot - m)|_{H^\rho(0,1)} \\ &\lesssim 2^{(j-\ell)/2} 2^{(\rho-1)(j-\ell)}, \end{aligned}$$

whence the result.

For a large class of elliptic differential/integral/pseudo-differential operators, the following estimate holds

$$2^{-r(|\lambda|+|\mu|)} |\langle A\psi_\mu, \psi_\lambda \rangle| \lesssim \frac{2^{-\beta ||\lambda|-|\mu||}}{[1 + d(\lambda, \mu)]^\gamma}$$

with

$$d(\lambda, \mu) = 2^{\min(|\lambda|, |\mu|)} \text{dist}(\text{supp } \psi_\lambda, \text{supp } \psi_\mu),$$

and

$\beta > n/2$  depends on the smoothness of the wavelets,

$\gamma > n$  is related to  $\tilde{L}$  and the order of the operator  $A$ .

These estimates allow an a-priori thresholding of the matrix elements, so that only 'non-negligible' elements are actually computed.

$\Rightarrow$  Savings in the construction of the matrix and the matrix-vector multiply.

Remark: Recent results on efficient approximate evaluation of matrix coefficients exploiting tensor-products, refinability, moment conditions, quadratures.

See [Dahmen - Schneider - Xu '98], [Bertoluzza - C. - Urban '99], [Berrone - Urban '99], [Dahmen - Schneider '99]

## Adaptivity

Assume that  $A : V \rightarrow V'$  is an algebraic and topological isomorphism,

$$\|Av\|_{V'} \asymp \|v\|_V, \quad \forall v \in V.$$

Let  $\Psi = \{\psi_\lambda\}_{\lambda \in \mathcal{M}}$  be a Riesz basis in  $V$ , and let  $\tilde{\Psi} = \{\tilde{\psi}_\lambda\}_{\lambda \in \mathcal{M}}$  be the dual biorthogonal basis in  $V'$ .

Exact Problem: Given  $f \in V'$ , find  $u \in V$  such that

$$\langle Au, v \rangle = \langle f, v \rangle, \quad \forall v \in V.$$

Galerkin Problem: Given  $\Lambda \subset \mathcal{M}$  and  $V_\Lambda = \text{span}\{\psi_\lambda : \lambda \in \Lambda\}$ , find  $u_\Lambda \in V_\Lambda$  such that

$$\langle Au_\Lambda, v_\Lambda \rangle = \langle f, v_\Lambda \rangle, \quad \forall v_\Lambda \in V_\Lambda.$$

Goal (fixed tolerance): Given  $\eta > 0$ , find in the most efficient way the smallest index set  $\Lambda \subset \mathcal{M}$  such that

$$\|u - u_\Lambda\|_V \sim \eta.$$

Goal (fixed resources): Given  $N > 0$ , find in the most efficient way an index set  $\Lambda \subset \mathcal{M}$  with  $\text{card } \Lambda = N$  such that

$$\|u - u_\Lambda\|_V \text{ is minimal.}$$

Precisely,

$$\|u - u_\Lambda\|_V \sim \inf_{v_N \in S_N} \|u - v_N\|_V$$

the cost of solving the problem is  $\mathcal{O}(N)$ .

This is a best N-term approximation problem, but  $u$  is not known!

Crucial step of any adaptive algorithm: Given the old set of active indices  $\Lambda \subset \mathcal{M}$  and the corresponding discrete solution  $u_\Lambda$ , define a new set  $\Lambda^* \subset \mathcal{M}$  such that

$$\|u - u_{\Lambda^*}\|_V < \|u - u_\Lambda\|_V.$$

The crucial step requires a *local error indicator*:

- inspect the wavelet coefficients of  $u_\Lambda$  (solution analysis)
- inspect the wavelet coefficients of the residual  $r_\Lambda := f - Au_\Lambda$  (residual analysis).

## Solution Analysis

[Liandrat - Tchamitchan '90, Maday - Perrier - Ravel '92, Bertoluzza - Maday - Ravel '94, C. - Cravero '97, Amat - Arandiga - Cohen - Donat '01]

Rationale: the wavelet coefficients of  $u$  are indicators of *local smoothness* “ $\Rightarrow$ ” the wavelet coefficients of  $u$  are indicators of potential *local error*.

A model strategy to define the new  $\Lambda^*$  is as follows:

Fix  $\varepsilon_{\max} > \varepsilon_{\min} > 0$ . For any  $\lambda \in \Lambda$ , look at  $(u_\Lambda)_\lambda$ :

$$|(u_\Lambda)_\lambda| \begin{cases} < \varepsilon_{\min} & \Rightarrow \lambda \notin \Lambda^* \\ \in [\varepsilon_{\min}, \varepsilon_{\max}] & \Rightarrow \lambda \in \Lambda^* \\ > \varepsilon_{\max} & \Rightarrow \lambda' \in \Lambda^* \text{ for all } \lambda' \text{ 'near' } \lambda. \end{cases}$$

### Pros

- Very easy and cheap to implement, particularly for evolution problems:

$$u_t + Au = f \quad \Rightarrow \quad u^{n+1} = u^n + \Delta t \Phi(u^n, \Delta t).$$

- Very effective when it works.

### Cons

- Mostly heuristic
- Theory is lacking, or partial
- The method may miss significant structures in the solution.

## Residual analysis

[Bertoluzza '95, Dahlke - Dahmen - Hochmuth - Schneider '97, Cohen - Masson '98, Cohen - Dahmen - DeVore '98, '00, Dahlke - Dahmen - Urban '01, Bertoluzza - Verani '01, C.- Urban '02]

Recall

$$\|v\|_V \asymp \|Av\|_{V'}, \quad \forall v \in V.$$

This implies

$$\|u - u_\Lambda\|_V \asymp \|Au - Au_\Lambda\|_{V'} \asymp \|f - Au_\Lambda\|_{V'}.$$

The quantity

$$r_\Lambda := f - Au_\Lambda$$

is the residual generated by the approximation  $u_\Lambda$  to  $u$ .

• **Key point:** Using Riesz bases  $\Psi$  in  $V$  and  $\tilde{\Psi}$  in  $V'$ , everything can be expressed in terms of sequences in  $\ell^2(\mathcal{M})$ :

$$u = \mathbf{u}^T \Psi = \sum_{\lambda \in \mathcal{M}} v_\lambda \psi_\lambda, \quad u_\Lambda = \mathbf{u}_\Lambda^T \Psi,$$

$$Au = (\mathbf{A}\mathbf{u})^T \tilde{\Psi}, \quad f = \mathbf{f}^T \tilde{\Psi}$$

with

$$\|u\|_V \asymp \|\mathbf{u}\|_{\ell^2(\mathcal{M})}, \quad \|f\|_{V'} \asymp \|\mathbf{f}\|_{\ell^2(\mathcal{M})},$$

$\mathbf{A} : \ell^2(\mathcal{M}) \rightarrow \ell^2(\mathcal{M})$  isomorphism.

Then,

$$\|\mathbf{u} - \mathbf{u}_\Lambda\|_{\ell^2(\mathcal{M})} \asymp \|\mathbf{f} - \mathbf{A}\mathbf{u}_\Lambda\|_{\ell^2(\mathcal{M})}.$$

From now on, we follow *Cohen - Dahmen - DeVore '00*. For simplicity, assume that  $\mathbf{A}$  is self-adjoint and coercive. Then

$$\begin{aligned} \mathbf{A} &\text{ is symmetric positive -- definite,} \\ \text{cond}_2(\mathbf{A}) &\asymp 1. \end{aligned}$$

**Basic Algorithm.** Fix a target accuracy  $\eta_T > 0$ . Approximate  $u$  by Richardson iterations

$$\begin{cases} \mathbf{u}^0 = \mathbf{0} \\ \mathbf{u}^{k+1} = \mathbf{u}^k + \alpha(\mathbf{f} - \mathbf{A}\mathbf{u}^k), \quad k \geq 0, \end{cases}$$

until

$$\|\mathbf{u} - \mathbf{u}^k\|_{\ell^2(\mathcal{M})} \asymp \eta_T.$$

Recall:  $\exists$  a range of  $\alpha > 0$  for which

$$\rho := \|\mathbf{I} - \alpha\mathbf{A}\|_{\ell^2 \rightarrow \ell^2} < 1,$$

so that the algorithm converges. Let us fix one of such  $\alpha$ .

**Definition.** For any  $\mathbf{v} \in \ell^2(\mathcal{M})$ , set

$$\Lambda = \text{supp } \mathbf{v} = \{\lambda \in \mathcal{M} : v_\lambda \neq 0\}.$$

The vector  $\mathbf{v}$  is finite if

$$\text{card } \Lambda < +\infty.$$

A finite vector with support  $\Lambda$  will be denoted by  $\mathbf{v}_\Lambda$ .

**Modified Algorithm.**

$$\begin{cases} \mathbf{u}_\Lambda^0 = \mathbf{0} \\ \mathbf{u}_\Lambda^{k+1} = (\mathbf{u}_\Lambda^k + \alpha(\mathbf{f}_\Lambda - (\mathbf{A}\mathbf{u}_\Lambda^k)_\Lambda))_\Lambda, \quad k \geq 0. \end{cases}$$

Precisely, suppose that we are able to construct finite vectors

$$\mathbf{f}_\Lambda := \text{RHS}[\mathbf{f}, \eta] \quad \text{such that} \quad \|\mathbf{f} - \mathbf{f}_\Lambda\|_{\ell^2(\mathcal{M})} \leq \eta,$$

$$\mathbf{w}_\Lambda := \text{MULT}[\mathbf{A}, \mathbf{v}_\Lambda, \eta] \quad \text{such that} \quad \|\mathbf{A}\mathbf{v}_\Lambda - \mathbf{w}_\Lambda\|_{\ell^2(\mathcal{M})} \leq \eta,$$

$$\bar{\mathbf{v}}_\Lambda := \text{COARSE}[\mathbf{v}_\Lambda, \eta] \quad \text{such that} \quad \|\mathbf{v}_\Lambda - \bar{\mathbf{v}}_\Lambda\|_{\ell^2(\mathcal{M})} \leq \eta$$

with minimal memory/complexity.

**Adaptive Algorithm.** Let  $K$  be a fixed integer.

i) Set  $\mathbf{u}_\Lambda^0 := \mathbf{0}$  and  $\eta_0 \sim \|\mathbf{f}\|_{\ell^2(\mathcal{M})}$

ii) for  $\ell \geq 0$  until  $\eta_\ell \leq \eta_T$  do:

a) Set  $\mathbf{v}_\Lambda^0 := \mathbf{u}_\Lambda^\ell$

b) for  $k = 0, \dots, K$  do:

compute  $\mathbf{f}_\Lambda^k = \text{RHS}[\mathbf{f}, \rho^k \eta_\ell]$  and  $\mathbf{w}_\Lambda^k := \text{MULT}[\mathbf{A}, \mathbf{v}_\Lambda^k, \rho^k \eta_\ell]$

update  $\mathbf{v}_\Lambda^{k+1} = \mathbf{v}_\Lambda^k + \alpha(\mathbf{f}_\Lambda^k - \mathbf{w}_\Lambda^k)$

c) set  $\mathbf{u}_\Lambda^{\ell+1} := \text{COARSE}[\mathbf{v}_\Lambda^{K+1}, 2\eta_\ell/5]$  and  $\eta_{\ell+1} := \eta_\ell/2$ .

**Proposition 1.** *There exists  $K > 0$  such that*

$$\|\mathbf{u} - \mathbf{v}_\Lambda^{K+1}\|_{\ell^2(\mathcal{M})} \leq \frac{1}{10}\eta^\ell.$$

**Corollary 2.** *The Adaptive Algorithm converges. Indeed, it is easily seen that*

$$\|\mathbf{u} - \mathbf{u}_\Lambda^{\ell+1}\|_{\ell^2(\mathcal{M})} \leq \eta_{\ell+1}.$$

**Question:** Is the Algorithm efficient?

Yes, because RHS, MULT and COARSE can be constructed taking into account the results of Nonlinear Approximation. The conclusion is as follows:

**Theorem 3.** *Let us assume that the exact solution  $u \in V$  is such that  $\mathbf{u} \in \ell_w^\tau(\mathcal{M})$ . Then*

$$\text{card}(\text{supp } \mathbf{u}_\Lambda^{\ell+1}) \lesssim \eta_{\ell+1}^{-d/s} \|\mathbf{u}\|_{\ell_w^\tau(\mathcal{M})}^{d/s}$$

and

$$\# \text{ floating point operations to compute } \mathbf{u}_\Lambda^{\ell+1} \lesssim \eta_{\ell+1}^{-d/s} \|\mathbf{u}\|_{\ell_w^\tau(\mathcal{M})}^{d/s}.$$

This means that the function  $u^{\ell+1}$  whose vector of coefficients is  $\mathbf{u}_\Lambda^{\ell+1}$  stays asymptotically close to the best  $N$ -term approximation of  $u$ . Furthermore, the operations needed to compute it is proportional to the number of its coefficients.